

# Assignment 5

March 1, 2017

1. Suppose  $X, \dots, X_n \stackrel{iid}{\sim} f_\theta(x)$  and let  $\beta$  denotes the power of the most powerful level  $\alpha$  test for testing  $H_0 : \theta = 0$  vs.  $H_1 : \theta = 1$ . Show that  $\alpha \leq \beta$ . Use the same logic to conclude that if  $\phi$  is a UMP level  $\alpha$  test then  $\phi$  is unbiased.
2. Suppose  $X$  has a distribution  $P_\theta$  for some  $\theta \in \Omega$ , and the null hypothesis  $H_0 : \theta \in \Omega_H$ . Assume that we have a test whose rejection region satisfy  $\mathcal{R}_\alpha \subset \mathcal{R}_{\alpha'}$  for  $\alpha < \alpha'$ .
  - (a) Show that for all  $\theta \in \Omega_H$ ,  $P_\theta(p(X) \leq u) \leq u$  for all  $0 \leq u \leq 1$ .
  - (b) If, for  $\theta \in \Omega_H$ ,  $P_\theta(X \in \mathcal{R}_\alpha) = \alpha$  for all  $0 \leq \alpha \leq 1$ , then  $P_\theta(p(\mathbf{X}) \leq u) = u$  for all  $0 \leq u \leq 1$ .
3. Let  $X_1, \dots, X_n \stackrel{iid}{\sim} Pois(\lambda)$ . Find the UMP level  $\alpha = 1 - e^{-n\lambda_0}(1 + n\lambda_0)$  test for  $H_0 : \lambda \leq \lambda_0$  vs.  $H_1 : \lambda > \lambda_0$ .
4. Let  $X_1, \dots, X_n \stackrel{iid}{\sim} U(0, \theta)$ . Find the UMP level  $\alpha$  test for testing  $H_0 : \theta = \theta_0$  vs.  $H_1 : \theta \neq \theta_0$ .
5. Let  $X_1, X_2$  be iid  $U(\theta, \theta + 1)$ . For testing  $H_0 : \theta = 0$  vs.  $H_1 : \theta > 0$ , we have two competing tests,

$$\phi_1(X_1) = 1 \text{ if } X_1 > .95; = 0 \text{ o.w.}$$

$$\phi_2(X_1, X_2) = 1 \text{ if } X_1 + X_2 > C; = 0 \text{ o.w.}$$

- (a) Find the value of  $C$  so that  $\phi_2$  has the same size as  $\phi_1$ .
- (b) Calculate the power function of each test.
- (c) Show how to get a test that has the same size but is more powerful than  $\phi_2$ .
6. Show that for a random sample  $X_1, \dots, X_n \stackrel{iid}{\sim} N(0, \sigma^2)$ . Find most powerful level  $\alpha$  test of  $H_0 : \sigma = \sigma_0$  vs.  $H_1 : \sigma = \sigma_1$ , where  $\sigma_0 < \sigma_1$ .
7. Suppose  $X_1, \dots, X_n$  are iid with a  $Beta(\mu, 1)$  and  $Y_1, \dots, Y_m$  are iid with a  $Beta(\theta, 1)$ . Also assume that  $X$ s are independent of  $Y$ s.
- (a) Find an LRT of  $H_0 : \theta = \mu$  vs.  $H_1 : \theta \neq \mu$ .
- (b) Show that the test above can be based on the statistic  $T = \frac{\sum_{i=1}^n \log(X_i)}{\sum_{i=1}^n \log(X_i) + \sum_{i=1}^m \log(Y_i)}$ .
8. Suppose that we have two independent random samples:  $X_1, \dots, X_n$  are  $Exp(\theta)$  and  $Y_1, \dots, Y_m$  are  $Exp(\mu)$ .
- (a) Find the LRT of  $H_0 : \theta = \mu$  vs.  $H_1 : \theta \neq \mu$ .
- (b) Show that the test above can be based on the statistic  $T = \frac{\sum_{i=1}^n X_i}{\sum_{i=1}^n X_i + \sum_{i=1}^m Y_i}$ .
- (c) Find the distribution of  $T$  when  $H_0$  is true.
9. Let  $X_1, \dots, X_n$  be a random sample from a  $N(\mu_X, \sigma_X^2)$ , and let  $Y_1, \dots, Y_m$  be an independent random sample from a  $N(\mu_Y, \sigma_Y^2)$ . We are interested in testing  $H_0 : \mu_X = \mu_Y$  vs.  $H_1 : \mu_X \neq \mu_Y$ , with the assumption that  $\sigma_X^2 = \sigma_Y^2$ . Derive the LRT test for this hypothesis. Provide an LRT test of size  $\alpha$ .
10. Let  $X_1, \dots, X_n$  be a random sample from a  $N(\theta, \sigma^2)$  population. Consider testing  $H_0 : \theta \leq \theta_0$  vs.  $H_1 : \theta > \theta_0$ . Let  $\bar{X}_m$  denote the sample mean of the first  $m$  observations,  $X_1, \dots, X_m$ , for  $m = 1, \dots, n$ . The test that rejects  $H_0$  when  $\bar{X}_m > \theta_0 + z_\alpha \sqrt{\sigma^2/m}$  is an unbiased size  $\alpha$  test.
11. In each of the following situations, calculate the p value of the observed data.

- (a) For testing  $H_0 : \theta \leq 0.5$  vs.  $H_1 : \theta > 0.5$ , 7 successes are observed out of 10 Bernoulli trials.
- (b) For testing  $H_0 : \lambda \leq 1$  vs.  $H_1 : \lambda > 1$ ,  $X = 3$  is observed, where  $X \sim Pois(\lambda)$ .

12. Let  $X$  be one observation from a Cauchy( $\theta$ ) distribution.

- (a) Show that the family does not have an MLR.
- (b) Show that the test

$$\phi(x) = \begin{cases} 1 & \text{if } 1 < X < 3 \\ 0 & \text{o.w.} \end{cases}$$

is most powerful of its size for testing  $H_0 : \theta = 0$  vs.  $H_1 : \theta = 1$ . Calculate the Type I and Type II error probabilities.

- (c) Is it a UMP test for testing  $H_0 : \theta = 0$  vs.  $H_1 : \theta > 0$ ? Prove or disprove.